

COMBINATORIAL RESULTS FOR CERTAIN SEMIGROUPS OF ORDER-PRESERVING FULL CONTRACTION MAPPINGS OF A FINITE CHAIN

A. D. Adeshola and A. Umar

Department of Mathematics, Kwara State University, Malete, Ilorin, Nigeria
Department of **Mathematical Sciences**, Petroleum Institute,
Khalifa University of Science and Technology, Abu Dhabi, U. A. E.
adeshola.dauda@kwasu.edu.ng and aumar@pi.ac.ae

Abstract

Let $\mathcal{T}_n$ be the full symmetric semigroup on $X_n = \{1, 2, \dots, n\}$ and let $\mathcal{OCT}_n$ and $\mathcal{ORCT}_n$ be its subsemigroups of order-preserving and order-preserving or order-reversing full contraction mappings of X_n , respectively. In this paper we investigate the cardinalities of some equivalences on $\mathcal{OCT}_n$ and $\mathcal{ORCT}_n$ which lead naturally to obtaining the orders of these subsemigroups.^{1, 2}

1 Introduction and Preliminaries

Let $X_n = \{1, 2, \dots, n\}$. A (partial) transformation $\alpha : \text{Dom } \alpha \subseteq X_n \rightarrow \text{Im } \alpha \subseteq X_n$ is said to be *full* or *total* if $\text{Dom } \alpha = X_n$; otherwise it is called *strictly* partial. The set of full transformations of X_n , denoted by $\mathcal{T}_n$, more commonly known as the *full transformation semigroup* is also known as the *full symmetric semigroup or monoid* with *composition of mappings* as the semigroup operation. We shall write $x\alpha$ for the image of x under α instead of $\alpha(x)$.

¹*Key Words*: height, right (left) waist and fix of a transformation, idempotents.

²Financial support from TRC Grant No.: RC/SCI/DOMS/13/01 is gratefully acknowledged.

The importance of $\mathcal{T}_n$ to semigroup theory may be likened to that of the symmetric group $\mathcal{S}_n$ to group theory. Every finite semigroup S is embeddable in $\mathcal{T}_n$ for some n , the analogue of Cayley's theorem for finite groups. Thus, just as the study of symmetric, alternating and dihedral groups has made a significant contribution to group theory, so has the study of various subsemigroups of $\mathcal{T}_n$. A transformation $\alpha \in \mathcal{T}_n$ is said to be *order-preserving* (order-reversing) if $(\forall x, y \in \text{Dom } \alpha) \ x \leq y \implies x\alpha \leq y\alpha$ ($x\alpha \geq y\alpha$) and, a *contraction mapping* (or simply a *contraction*) if $(\forall x, y \in \text{Dom } \alpha) \ |x - y| \geq |x\alpha - y\alpha|$. We shall denote by $\mathcal{OCT}_n$ and $\mathcal{ORCT}_n$, the semigroups of order-preserving full contractions and of order-preserving or order-reversing full contractions of X_n , respectively. We shall also denote by $\mathcal{ODCT}_n$, the semigroup of order-preserving and order-decreasing full contractions of X_n .

Recently, Zhao and Yang [20] initiated the algebraic study of semigroups of order-preserving partial contractions of X_n , where they referred to our contractions as *compressions*. Earlier one of the authors [16] studied $\mathcal{DT}_n$, the semigroup of order-decreasing full transformations of X_n . This paper investigates the combinatorial properties of $\mathcal{OCT}_n$, $\mathcal{ORCT}_n$ and $\mathcal{ODCT}_n$, and of their subsets of idempotents.

In this section we introduce basic terminologies and prove some preliminary results. In Section 2 we obtain the cardinalities of various equivalence classes defined on $\mathcal{OCT}_n$ and in Sections 3 and 4 we obtain the analogues of the results in Section 2 for $\mathcal{ORCT}_n$ and $\mathcal{ODCT}_n$, respectively. These cardinalities lead to formulae for the orders of $\mathcal{OCT}_n$, $\mathcal{ORCT}_n$ and $\mathcal{ODCT}_n$ as well as new triangles of numbers that as a result of this work were recently recorded in [15].

For standard concepts in semigroup and transformation semigroup theory, see for example [7, 3]. In particular, the set of idempotents of a semigroup S will be denoted by $E(S)$.

Let

$$\mathcal{OR}_n = \{\alpha \in \mathcal{T}_n : (\forall x, y \in X_n) \ x \leq y \implies x\alpha \leq y\alpha \text{ or } x\alpha \geq y\alpha\} \quad (1)$$

be the subsemigroup of $\mathcal{T}_n$ consisting of all order-preserving or order-reversing full transformations of X_n , and let

$$\mathcal{O}_n = \{\alpha \in \mathcal{T}_n : (\forall x, y \in X_n) \ x \leq y \implies x\alpha \leq y\alpha\} \quad (2)$$

be the subsemigroup of $\mathcal{T}_n$ consisting of all order-preserving full transformations of X_n . Also let

$$\mathcal{CT}_n = \{\alpha \in \mathcal{T}_n : (\forall x, y \in X_n) \ |x - y| \geq |x\alpha - y\alpha|\} \quad (3)$$

be the subsemigroup of $\mathcal{T}_n$ consisting of all full contractions of X_n , and let

$$\mathcal{D}_n = \{\alpha \in \mathcal{T}_n : (\forall x \in X_n) \ x\alpha \leq x\} \quad (4)$$

be the subsemigroup of $\mathcal{T}_n$ consisting of all *order-decreasing* full transformations of X_n .

The *height* of a partial transformation α is denoted and defined by $h(\alpha) = |\text{Im } \alpha|$, the *right [left] waist* of α is denoted and defined by $w^+(\alpha) = \max(\text{Im } \alpha)$ [$w^-(\alpha) = \min(\text{Im } \alpha)$], the *fix* of α is denoted and defined by $f(\alpha) = |F(\alpha)|$, where

$$F(\alpha) = \{x \in \text{Dom } \alpha : x\alpha = x\}.$$

It is also well-known that a partial transformation ϵ is *idempotent* ($\epsilon^2 = \epsilon$) if and only if $\text{Im } \epsilon = F(\epsilon)$. It is worth noting that to define the left (right) waist of a transformation the base set X_n must be totally ordered.

We have the following results:

Lemma 1.1 *Let $\alpha \in \mathcal{CT}_n$ be such that $f(\alpha) = m$. Then $F(\alpha) = \{i, i+1, \dots, i+m-1\}$. Equivalently, $F(\alpha)$ is convex.*

Proof. Observe that it is sufficient to show that any point between two fixed points (of $\alpha \in \mathcal{CT}_n$) must also be a fixed point. Let $x, y \in F(\alpha)$. Then $x\alpha = x$ and $y\alpha = y$. Suppose also without loss of generality $x < x' < y$ for some $x' \in X_n$. Note that if $x' = x'\alpha$, there is nothing to prove. Thus we consider two cases: (i) $x' > x'\alpha$; (ii) $x' < x'\alpha$. In the former, we have

$$|y - x'| = |y\alpha - x'| = y\alpha - x' < y\alpha - x'\alpha = |y\alpha - x'\alpha|$$

which implies that α is not a contraction. Hence we get a contradiction. In the latter, we have

$$|x' - x| = |x' - x\alpha| = x' - x\alpha < x'\alpha - x\alpha = |x'\alpha - x\alpha|$$

which implies that α is not a contraction. Hence we get a contradiction. Thus, the proof is complete. $\square$

Lemma 1.2 *Let $\alpha \in \mathcal{CT}_n$ be such that $h(\alpha) = p$. Then $\text{Im } \alpha = \{i, i+1, \dots, i+p-1\}$. Equivalently, $\text{Im } \alpha$ is convex.*

Proof. (By contradiction) Suppose that $\text{Im } \alpha$ is not convex. Then there exist $x, z \in \text{Im } \alpha$ with $x < y < z$ for some $y \in X_n \setminus \text{Im } \alpha$. Let $(y-1]$ and $[y+1)$ be the lower and upper saturations of $y-1$ and $y+1$, respectively. Notice that $x \in (y-1]$ and $z \in [y+1)$. Moreover, $(y-1]\alpha^{-1} \neq X_n \neq [y+1)\alpha^{-1}$ but $(y-1]\alpha^{-1} \cup [y+1)\alpha^{-1} = X_n$. If $(y-1]\alpha^{-1}$ is convex then since

$(y-1]\alpha^{-1} \neq X_n$ there exist either (i) $t \in (y-1]\alpha^{-1}$ and $t+1 \in [y+1)\alpha^{-1}$; or (ii) $t \in (y-1]\alpha^{-1}$ and $t-1 \in [y+1)\alpha^{-1}$.

Case (i): it is clear that $t\alpha \leq y-1$ and $(t+1)\alpha \geq y+1$ so that

$$2 \leq (t+1)\alpha - t\alpha = |(t+1)\alpha - t\alpha| \leq |(t+1) - t| = 1,$$

which is a contradiction.

Case (ii): it is clear that $t\alpha \leq y-1$ and $(t-1)\alpha \geq y+1$ so that

$$2 \leq (t-1)\alpha - t\alpha = |(t-1)\alpha - t\alpha| \leq |(t-1) - t| = 1,$$

which is another contradiction.

On the other hand if $(y-1]\alpha^{-1}$ is not convex then there exists $t \in (y-1]\alpha^{-1}$ and either $t+1 \in [y+1)\alpha^{-1}$ or $t-1 \in [y+1)\alpha^{-1}$. In the former, we see that $t\alpha \leq y-1$ and $(t+1)\alpha \geq y+1$. Thus,

$$2 \leq (t+1)\alpha - t\alpha = |(t+1)\alpha - t\alpha| \leq |(t+1) - t| = 1,$$

which is a contradiction. In the latter, we see that $t\alpha \leq y-1$ and $(t-1)\alpha \geq y+1$. Thus,

$$2 \leq (t-1)\alpha - t\alpha = |(t-1)\alpha - t\alpha| \leq |(t-1) - t| = 1,$$

which is another contradiction. Hence, the proof is complete. $\square$

Next we state three important identities that will be needed later.

Lemma 1.3 (*Vandemonde's Convolution Identity, [14, (3a), p.8]*). For all natural numbers m, n and p we have

$$\sum_{k=0}^n \binom{n}{m-k} \binom{p}{k} = \binom{n+p}{m}.$$

Lemma 1.4 [*11, Lemma 1.3*] For all natural numbers j and a we have

$$\sum_{i=0}^{j-a} \binom{j-i}{a} = \binom{j+1}{a+1}.$$

Lemma 1.5 ([*14, (3b), p.8*]). For all natural numbers m, n and p we have

$$\sum_{k=0}^n \binom{n-k}{n-m} \binom{p+k-1}{k} = \binom{n+p}{m}.$$

2 Order-preserving Full Contractions

Enumerative problems of an essentially combinatorial nature arise naturally in the study of semigroups of transformations. Many numbers and triangles of numbers regarded as combinatorial gems like the Stirling numbers [6, pp. 42 & 96], the factorial [16], the Fibonacci number [5], Catalan numbers [3], Lah numbers [3], etc., have all featured in these enumeration problems. For a nice survey article concerning combinatorial problems in the partial transformation semigroup and some of its subsemigroups we refer the reader to Umar [18]. These enumeration problems lead to many numbers in Sloane's encyclopaedia of integer sequences [15] but there are also others that are not yet or have just been recorded in [15].

Now recall the definitions of *height*, (*right*) *waist* and *fix* of $\alpha \in \mathcal{T}_n$ stated earlier. From Umar [18], we quote this result.

Lemma 2.1 *Let $X_n = \{1, 2, \dots, n\}$ and $P = \{p, m, k\}$, where for a given $\alpha \in \mathcal{T}_n$, we set $p = h(\alpha)$, $m = f(\alpha)$ and $k = w^+(\alpha)$. Then we have the following:*

1. $n \geq k \geq p \geq m \geq 1$;
2. $k = 1 \implies p = 1 \implies m \leq 1$.

As in Umar [18] let S be a set of full transformations on X_n and consider the combinatorial functions:

$$\begin{aligned}
 F(n; p, m, k) &= |\{\alpha \in S : \wedge(h(\alpha) = p, f(\alpha) = m, w^+(\alpha) = k)\}|, \\
 F(n; p, m) &= |\{\alpha \in S : \wedge(h(\alpha) = p, f(\alpha) = m)\}|, \\
 F(n; p, k) &= |\{\alpha \in S : \wedge(h(\alpha) = p, w^+(\alpha) = k)\}|, \\
 F(n; m, k) &= |\{\alpha \in S : \wedge(f(\alpha) = m, w^+(\alpha) = k)\}|, \\
 F(n; p) &= |\{\alpha \in S : h(\alpha) = p\}|, \\
 F(n; m) &= |\{\alpha \in S : f(\alpha) = m\}|, \\
 F(n; k) &= |\{\alpha \in S : w^+(\alpha) = k\}|.
 \end{aligned}$$

Observe that

$$F(n; a_1, a_2) = \sum_{a_3} F(n; a_1, a_2, a_3), \quad F(n; a_1) = \sum_{a_2} F(n; a_1, a_2),$$

and

$$|S| = \sum_{a_1} F(n; a_1)$$

where $\{a_1, a_2, a_3\} = \{p, m, k\}$.

The following lemma is crucial to our investigation.

Lemma 2.2 *Let*

$$\begin{aligned} a_{n,p} &= |\{\alpha \in \mathcal{OCT}_n : (F(\alpha) = \{1\}) \wedge (h(\alpha) = p)\}| \\ &= |\{\alpha \in \mathcal{OCT}_n : (F(\alpha) = \{n\}) \wedge (h(\alpha) = p)\}|. \end{aligned}$$

Then $a_{n,p} = \binom{n-2}{p-1}$.

Proof. Let $\alpha \in \mathcal{OCT}_n$ be such that $F(\alpha) = \{1\}$ and $h(\alpha) = p$. First observe that $\text{Im } \alpha = \{1, 2, \dots, p\}$, by Lemma 1.2. Next, recall that the blocks $(y\alpha^{-1} \text{ for } y \in \text{Im } \alpha)$ are convex by order-preservedness and so the number of partitions of X_n into p convex blocks is the number of ways of inserting $p - 1$ bars between the $n - 1$ spaces (between the points of the n -chain). However, notice that in this case the space between 1 and 2 is not available so that 2 does not become a fixed point since $F(\alpha) = \{1\}$. Thus, we have $n - 2$ available places to insert $p - 1$ bars, which can be done in $\binom{n-2}{p-1}$ ways, without new fixed points being introduced. The result when $F(\alpha) = \{n\}$, follows by symmetry. $\square$

The main result in this paper can now be stated.

Proposition 2.3 *Let* $S = \mathcal{OCT}_n = \mathcal{O}_n \cap \mathcal{CT}_n$. *Then* $F(n; p, m, k) = \binom{n-m-1}{n-p-1}$, *for* $(n > p \geq m \geq 1)$.

Proof. Let $\alpha \in \mathcal{OCT}_n$ be such that $h(\alpha) = p$, $f(\alpha) = m$ and $w^+(\alpha) = k$. Then by Lemmas 1.1 & 1.2 we see that

$$F(\alpha) = \{i, i + 1, \dots, i + m - 1\} \subseteq \{k - p + 1, k - p + 2, \dots, k\} = \text{Im } \alpha,$$

where $k - p + 1 \leq i \leq k - m + 1$. Next, observe that by order-preservedness we may decompose α into $\beta = \alpha|_{\{1, 2, \dots, i\}}$, $id_{\{i+1, i+2, \dots, i+m-2\}}$ and $\beta' = \alpha|_{\{i+m-1, i+m, \dots, n\}}$, where $F(\beta) = \{i\}$, $\text{Im } \beta = \{k - p + 1, \dots, i\}$, $F(\beta') = \{i + m - 1\}$, and $\text{Im } \beta' = \{i + m - 1, \dots, k\}$. Now it is not difficult to see that for β there are $a_{i, i-k+p}$ possible maps, and for β' there are $a_{n-i-m+2, k-i-m+2}$ possible maps. Thus, multiplying the two numbers and

taking the sum of the product over i from $k - p + 1$ to $k - m + 1$ and using Lemmas 1.5 & 2.2 gives

$$\begin{aligned}
F(n; p, k, m) &= \sum_{i=k-p+1}^{k-m+1} a_{i, i-k+p} a_{n-i-m+2, k-i-m+2} \\
&= \sum_{i=k-p+1}^{k-m+1} \binom{i-2}{i-k+p-1} \binom{n-i-m}{k-i-m+1} \\
&= \sum_{i=k-p+1}^{k-m+1} \binom{i-2}{i-k+p-1} \binom{n-i-m}{n-k-1} \\
&= \binom{n-m-1}{p-m} = \binom{n-m-1}{n-p-1},
\end{aligned}$$

using the substitution $i - k + p - 1 = j$, for the last step. $\square$

Corollary 2.4 *Let $S = \mathcal{OCT}_n$. Then $F(n; p, m) = (n - p + 1) \binom{n-m-1}{n-p-1}$, for $n > p \geq m \geq 1$.*

Corollary 2.5 *Let $S = \mathcal{OCT}_n$. Then $F(n; p, k) = \binom{n-1}{p-1}$, for $k \geq p \geq 1$.*

Proof. It follows directly from Proposition 2.3 and Lemma 1.4. $\square$

Corollary 2.6 *Let $S = \mathcal{OCT}_n$. Then $F(n; m, k) = \sum_{p=m}^k \binom{n-m-1}{n-p-1}$, for $k \geq m \geq 1$.*

Proof. It follows directly from Proposition 2.3. $\square$

Corollary 2.7 *Let $S = \mathcal{OCT}_n$. Then $F(n; p) = (n - p + 1) \binom{n-1}{p-1}$, for $p \geq 1$.*

Proof. It follows from either of the Corollaries 2.4 & 2.5. $\square$

Corollary 2.8 *Let $S = \mathcal{OCT}_n$. Then $F(n; k) = \sum_{p=1}^k \binom{n-1}{p-1}$, for $k \geq 1$.*

Proof. It follows from either of the Corollaries 2.5 & 2.6. $\square$

For $0 \leq i \leq n$, let

$$F(n; m_i) = | \{ \alpha \in S : f(\alpha) = i \} |.$$

Corollary 2.9 Let $S = \mathcal{OCT}_n$. Then $F(n; m_n) = 1$ and $F(n; m) = (n - m + 3)2^{n-m-2}$, for $n \geq 2$ and $n > m \geq 1$.

Proof. $F(n; m_n) = 1$ follows from the fact that the identity transformation is the only transformation that has n fixed points. Next observing that $F(n; p = n, m < n) = 0$ and using Corollary 2.4, we see that

$$\begin{aligned}
F(n; m) &= \sum_{p=m}^n F(n; p, m) = \sum_{p=m}^{n-1} F(n; p, m) + F(n; p = n, m < n) \\
&= \sum_{p=m}^{n-1} (n - p + 1) \binom{n - m - 1}{n - p - 1} \\
&= \sum_{p=m}^{n-1} (n - p - 1) \binom{n - m - 1}{n - p - 1} + 2 \sum_{p=m}^{n-1} \binom{n - m - 1}{n - p - 1} \\
&= (n - m - 1) \sum_{p=m}^{n-1} \binom{n - m - 2}{n - p - 2} + 2(2^{n-m-1}) \\
&= (n - m - 1)2^{n-m-2} + 2^2(2^{n-m-2}) = (n - m + 3)2^{n-m-2},
\end{aligned}$$

as required. $\square$

Corollary 2.10 Let $S = \mathcal{OCT}_n$. Then $|S| = |\mathcal{OCT}_n| = (n + 1)2^{n-2}$, for $n \geq 1$.

Proof. It follows from any one of the Corollaries 2.7, 2.8 & 2.9. $\square$

Corollary 2.11 Let $S = E(\mathcal{OCT}_n)$. Then $F(n; p, k) = F(n; m, k) = 1$, for $k \geq p = m \geq 1$.

Proof. Since $F(\alpha) = \text{Im } \alpha$ for idempotents, it follows that $p = m$. Hence the result follows from Proposition 2.3. $\square$

Corollary 2.12 Let $S = E(\mathcal{OCT}_n)$. Then $F(n; p) = F(n; m) = n - p + 1 = n - m + 1$, for $p = m \geq 1$.

Corollary 2.13 Let $S = E(\mathcal{OCT}_n)$. Then $F(n; k) = k$, for $k \geq 1$.

Corollary 2.14 $|E(\mathcal{OCT}_n)| = n(n + 1)/2 = \binom{n+1}{2}$, for $n \geq 1$.

Remark 2.15 The triangle of numbers $F(n; m)$ has as a result of this work been recorded in [15]. However, $F(n; p)$, $F(n; k)$, $F(n; m_1)$ and $|\mathcal{OCT}_n|$ were recorded (in [15]) as A104698, A008949, A045623 and A001792, respectively.

3 Order-preserving or Order-reversing Full Contractions

Remark 3.1 For $h(\alpha) = p = 1$ the concepts of order-preserving and order-reversing coincide but distinct otherwise. However, the map $\alpha \mapsto \alpha h$, where $xh = n - x + 1$, for all x in X_n is a bijection between the two sets for $p \geq 2$, see [2, page 2, last paragraph].

Remark 3.2 Every idempotent is necessarily order-preserving. Thus, there are no additional idempotents from reversing the order.

The main result of this section is

Proposition 3.3 Let $S = \mathcal{ORCT}_n = \mathcal{OR}_n \cap \mathcal{CT}_n$. Then

$$F(n; p, k) = \begin{cases} 2\binom{n-1}{p-1}, & p > 1; \\ 1, & \text{otherwise.} \end{cases}$$

Proof. It follows from Corollary 2.5 and Remark 3.1. □

Corollary 3.4 Let $S = \mathcal{ORCT}_n$. Then

$$F(n; p) = \begin{cases} 2(n - p + 1)\binom{n-1}{p-1}, & p > 1; \\ n, & \text{otherwise.} \end{cases}$$

Proof. It follows from Proposition 3.3 and the fact that $p \leq k \leq n$. □

Corollary 3.5 Let $S = \mathcal{ORCT}_n$. Then $F(n; k) = 2\sum_{p=1}^k \binom{n-1}{p-1} - 1$, for $k \geq 1$.

Proof. It follows from Proposition 3.3 and the fact that $1 \leq p \leq k$. □

Corollary 3.6 Let $S = \mathcal{ORCT}_n$. Then $|S| = |\mathcal{ORCT}_n| = (n+1)2^{n-1} - n$, for $n \geq 1$.

Proof. It follows from either of the Corollaries 3.4 & 3.5. $\square$

Remark 3.7 *The triangles of numbers $F(n; p)$, $F(n; k)$ and $F(n; m)$ have as a result of this work just been recorded in [15].*

4 Order-preserving and Order-decreasing Full Contractions

The main result of this section is

Proposition 4.1 *Let $S = \mathcal{ODCT}_n = \mathcal{D}_n \cap \mathcal{OCT}_n$. Then*

$$F(n; p, k, m) = \begin{cases} \binom{n-m-1}{p-m}, & p = k \text{ \& } n > m; \\ 0, & \text{otherwise.} \end{cases}$$

Proof. Let $\alpha \in \mathcal{ODCT}_n$ be such that $h(\alpha) = p$, $f(\alpha) = m$ and $w^+(\alpha) = k$. Then by Lemmas 1.1 & 1.2 we see that

$$F(\alpha) = \{1, 2, \dots, m\} \subseteq \{1, 2, \dots, p\} = \text{Im } \alpha.$$

Moreover, $p = k$. Next, observe that by order-preservedness we may decompose α into $id_{\{1, 2, \dots, m-1\}}$ and $\beta' = \alpha|_{\{m, m+1, \dots, n\}}$, where $F(\beta') = \{m\}$, and $\text{Im } \beta' = \{m, m+1, \dots, p\}$. Now it is not difficult to see that there are $a_{n-m+1, p-m+1}$ possible β' s, by Lemma 2.2. Thus,

$$F(n; p, k, m) = \begin{cases} \binom{n-m-1}{p-m}, & p = k \text{ \& } n > m; \\ 0, & \text{otherwise.} \end{cases} \quad \square$$

Corollary 4.2 *Let $S = \mathcal{ODCT}_n$. Then $F(n; p, m) = \binom{n-m-1}{p-m}$, for $p \geq m \geq 1$ & $n > m$.*

Corollary 4.3 *$S = \mathcal{ODCT}_n$. Then $F(n; m, k) = \binom{n-m-1}{k-m}$, for $k \geq m \geq 1$ & $n > m$.*

Corollary 4.4 *Let $S = \mathcal{ODCT}_n$. Then $F(n; p) = \binom{n-1}{p-1}$, for $p \geq 1$.*

Corollary 4.5 *Let $S = \mathcal{ODCT}_n$. Then $F(n; k) = \binom{n-1}{k-1}$, for $k \geq 1$.*

Corollary 4.6 *Let $S = \mathcal{ODCT}_n$. Then $F(n; m_n) = 1$ and $F(n; m) = 2^{n-m-1}$, for $n > m \geq 1$.*

Corollary 4.7 Let $S = \mathcal{ODCT}_n$. Then $|S| = |\mathcal{ODCT}_n| = 2^{n-1}$, for $n \geq 1$.

Corollary 4.8 Let $S = E(\mathcal{ODCT}_n)$. Then

$$F(n; p, m) = \begin{cases} 1, & p = m; \\ 0, & \text{otherwise.} \end{cases}$$

Proof. Since $F(\alpha) = \text{Im } \alpha$ for idempotents, it follows that $p = m$. Hence the result follows from Proposition 2.3. $\square$

Corollary 4.9 Let $S = E(\mathcal{ODCT}_n)$. Then $F(n; p) = F(n; m) = F(n; k) = 1$, for $p, k, m \geq 1$.

Corollary 4.10 $|E(\mathcal{ODCT}_n)| = n$, for $n \geq 1$.

Acknowledgements. The first named author would like to thank Kwara State University, Malete and TET Fund for financial support, and Sultan Qaboos University for hospitality.

References

- [1] BORWEIN, D., RANKIN, S. AND RENNER, L. Enumeration of injective partial transformations. *Discrete Math.* **73** (1989), 291–296.
- [2] FERNANDES, V. H., GOMES, G. M. S. AND JESUS, M. M. The cardinal and idempotent number of various monoids of transformations on a finite chain. *Bull. Malays. Math. Sci. Soc.* **34** (2011), 79–85.
- [3] GANYUSHKIN, O. AND MAZORCHUK, V. *Classical Finite Transformation Semigroups: An Introduction*. Springer, London, 2009.
- [4] HIGGINS, P. M. Combinatorial results for semigroups of order-preserving mappings. *Math. Proc. Camb. Phil. Soc.* **113** (1993), 281–296.
- [5] HOWIE, J. M. Products of idempotents in certain semigroups of transformations. *Proc. Edinburgh Math. Soc.* **17** (1971), 223–236.
- [6] HOWIE, J. M. Combinatorial and probabilistic results in transformation semigroups. *Words, languages and combinatorics. II (Kyoto, 1992)*, World Sci. Publ., River Edge, NJ, (1994), 200–206.

- [7] HOWIE, J. M. *Fundamentals of semigroup theory*. Oxford: Clarendon Press, 1995.
- [8] LALLEMENT, G. *Semigroups and combinatorial applications*. Wiley, New York 1979.
- [9] LARADJI, A. AND UMAR, A. On certain finite semigroups of order-decreasing transformations I. *Semigroup Forum* **69** (2004), 184–200.
- [10] LARADJI, A. AND UMAR, A. Combinatorial results for semigroups of order-preserving partial transformations. *Journal of Algebra* **278** (2004), 342–359.
- [11] LARADJI, A. AND UMAR, A. Combinatorial results for semigroups of order-decreasing partial transformations. *J. Integer Seq.* **7** (2004), Article 04.3.8.
- [12] LARADJI, A. AND UMAR, A. Combinatorial results for semigroups of order-preserving full transformations. *Semigroup Forum* **72** (2006), 51–62.
- [13] LARADJI, A. AND UMAR, A. Some combinatorial properties of the symmetric monoid. *Int. Journal of Alg. and Computations* **21** (2011), 857–865.
- [14] RIORDAN, J. *Combinatorial Identities*, John Wiley and Sons, New York, 1968.
- [15] SLOANE, N. J. A. (Ed.), *The On-Line Encyclopedia of Integer Sequences*, 2011. Available at <http://oeis.org/>.
- [16] UMAR, A. On the semigroups of order-decreasing finite full transformations. *Proc. Roy. Soc. Edinburgh* **120A** (1992), 129–142.
- [17] UMAR, A. Enumeration of certain finite semigroups of transformations. *Discrete Math.* **89** (1998), 291–297.
- [18] UMAR, A. Some combinatorial problems in the theory of partial transformation semigroups. *Algebra Discrete Math.* **17** (2014), 110–134.
- [19] UMAR, A. Combinatorial results for semigroups of orientation-preserving partial transformations. *J. Integer Seq.* **14** (2011), Article 11.7.5.
- [20] ZHAO, P. AND YANG, M. Regularity and Green’s relations on semigroups of transformations preserving order and compression. *Bull. Korean Math. Soc.* **49** (2012), 1015–1025.